

Boundary value problems

Example 147. The IVP (initial value problem) $y'' + 4y = 0$, $y(0) = 0$, $y'(0) = 0$ has the unique solution $y(x) = 0$.

Initial value problems are often used when the problem depends on time. Then, $y(0)$ and $y'(0)$ describe the initial configuration at $t = 0$.

For problems which instead depend on spatial variables, such as position, it may be natural to specify values at positions on the boundary (for instance, if $y(x)$ describes the steady-state temperature of a rod at position x , we might know the temperature at the two end points).

The next example illustrates that such a boundary value problem (BVP) may or may not have a unique solution.

Example 148. Verify the following claims.

- (a) The BVP $y'' + 4y = 0$, $y(0) = 0$, $y(1) = 0$ has the unique solution $y(x) = 0$.
- (b) The BVP $y'' + \pi^2 y = 0$, $y(0) = 0$, $y(1) = 0$ is solved by $y(x) = B \sin(\pi x)$ for any value B .

Solution.

- (a) We know that the general solution to the DE is $y(x) = A \cos(2x) + B \sin(2x)$. The boundary conditions imply $y(0) = A \stackrel{!}{=} 0$ and, using that $A = 0$, $y(1) = B \sin(2) \stackrel{!}{=} 0$ shows that $B = 0$ as well.
- (b) This time, the general solution to the DE is $y(x) = A \cos(\pi x) + B \sin(\pi x)$. The boundary conditions imply $y(0) = A \stackrel{!}{=} 0$ and, using that $A = 0$, $y(1) = B \sin(\pi) \stackrel{!}{=} 0$. This second condition is true for every B .

It is therefore natural to ask: for which λ does the BVP $y'' + \lambda y = 0$, $y(0) = 0$, $y(L) = 0$ have nonzero solutions? (We assume that $L > 0$.)

Such solutions are called **eigenfunctions** and λ is the corresponding **eigenvalue**.

Remark. Compare that to our previous use of the term eigenvalue: given a matrix A , we asked: for which λ does $Av - \lambda v = 0$ have nonzero solutions v ? Such solutions were called eigenvectors and λ was the corresponding eigenvalue.

Example 149. Find all eigenfunctions and eigenvalues of $y'' + \lambda y = 0$, $y(0) = 0$, $y(L) = 0$.

Such a problem is called an **eigenvalue problem**.

Solution. The solutions of the DE look different in the cases $\lambda < 0$, $\lambda = 0$, $\lambda > 0$, so we consider them individually.

$\lambda = 0$. Then $y(x) = Ax + B$ and $y(0) = y(L) = 0$ implies that $y(x) = 0$. No eigenfunction here.

$\lambda < 0$. The roots of the characteristic polynomial are $\pm\sqrt{-\lambda}$. Writing $\rho = \sqrt{-\lambda}$, the general solution therefore is $y(x) = Ae^{\rho x} + Be^{-\rho x}$. $y(0) = A + B \stackrel{!}{=} 0$ implies $B = -A$. Using that, we get $y(L) = A(e^{\rho L} - e^{-\rho L}) \stackrel{!}{=} 0$. For eigenfunctions we need $A \neq 0$, so $e^{\rho L} = e^{-\rho L}$ which implies $\rho L = -\rho L$. This cannot happen since $\rho \neq 0$ and $L \neq 0$. Again, no eigenfunctions in this case.

$\lambda > 0$. The roots of the characteristic polynomial are $\pm i\sqrt{\lambda}$. Writing $\rho = \sqrt{\lambda}$, the general solution thus is $y(x) = A \cos(\rho x) + B \sin(\rho x)$. $y(0) = A \stackrel{!}{=} 0$. Using that, $y(L) = B \sin(\rho L) \stackrel{!}{=} 0$. Since $B \neq 0$ for eigenfunctions, we need $\sin(\rho L) = 0$. This happens if $\rho L = n\pi$ for $n = 1, 2, 3, \dots$ (since ρ and L are both positive, n must be positive as well). Equivalently, $\rho = \frac{n\pi}{L}$. Consequently, we find the eigenfunctions $y_n(x) = \sin\left(\frac{n\pi x}{L}\right)$, $n = 1, 2, 3, \dots$, with eigenvalue $\lambda = \left(\frac{n\pi}{L}\right)^2$.